

SHORT COMMUNICATION

A lectotype in *Banksia* L.f. (Proteaceae) clarified

Recently I realised that I have lectotypified the name *Banksia* L.f. sect. *Oncostylis* Benth. (Proteaceae) twice, choosing different species. Bentham published the name in *Flora Australiensis* (Bentham 1870: 542, 544) and included 13 species in the section, where all remain in the current classification (George 1999). The protologue gives little lead in choosing a lectotype. In *Nuytsia* (George 1996: 21), I chose *B. ericifolia* L.f. as the 'earliest-described of the included species'. In *Flora of Australia* (George 1999: 397), I chose *B. sphaerocarpa* R.Br. as the 'first-described species with the leaf character states given first in the protologue'. Either species would be appropriate but *B. ericifolia* has been adopted for the Australian Plant Name Index (APNI 2012) and is here confirmed as lectotype of the name.

References

- APNI (2012). *Australian Plant Name Index*, IBIS database (Centre for Australian National Biodiversity Research, Australian Government, Canberra). <http://www.cpbr.gov.au/cgi-bin/apni> [accessed 5 September 2012].
- Bentham, G. (1870). *Flora Australiensis*. Vol. 5 (Reeve & Co.: London.)
- George, A.S. (1996). Notes on *Banksia* L.f. *Nuytsia* 11: 21–24.
- George, A.S. (1999). *Banksia*. In: Orchard, A.E., Thompson, H.S., & McCarthy, P.M. (eds) *Flora of Australia*. Vol. 17B, pp. 175–251. (Australian Biological Resources Study: Canberra, and CSIRO Publishing: Melbourne.)

Alex S. George

'Four Gables', 18 Barclay Road, Kardinya,
Western Australia 6163